arXiv:1605.03145v2 [cond-mat.mes-hall] 11 May 2016

Microscopic Friction Emulators

Davide Mandelli[1] & Erio Tosatti[1,2,3]

[1] *International School for Advanced Studies (SISSA), Via Bonomea 265, 34136 Trieste, Italy*
[2] *CNR-IOM Democritos National Simulation Center, Via Bonomea 265, 34136 Trieste, Italy*
[3] *International Centre for Theoretical Physics (ICTP), Strada Costiera 11, 34014 Trieste, Italy*

(Dated: May 12, 2016)

Cold ions sliding across periodic potential patterns formed by lasers elucidate the physics of dry friction between crystals. Experiments with no more than six ions suffice to explore a vast domain of frictional forces.

The study of friction is a centuries-old, but still vibrant, subject. The simplest, archetypal dry sliding friction problem is that between crystal surfaces, more precisely the force resisting the relative lateral motion of two crystals in contact. In this case, the periodic spacing of the atoms in the crystal eliminates the complexity introduced by the ill-defined nature of ordinary, non-crystalline surfaces. Studies of nanometre-scale systems through non-equilibrium theory, computer simulations, and most recently the use of artificial friction emulators have revived this topic.[1] Writing in Science and in Nature Physics, respectively, Bylinskii et al.[2] and Gangloff et al.[3] follow up on earlier theoretical suggestions,[4–6] and present friction emulators formed from trapped ions that slide over optical lattices; the latter are periodic intensity potentials created by the interference of laser beams.

Although dry sliding is a well-defined problem, sufficient inroads have not yet been made into it – microscopically well defined experimental systems are not abundant. Theory of frictional dynamics is often limited to computer simulations – vivid but often hasty and incomplete representations of the real, complex and slow sliding of true solid interfaces. Much of our understanding relies on apparently trivial one-dimensional periodic potential models, such as the Prandt-Tomlinson (PT) single-slider[7] and the Frenkel-Kontorova (FK) sliding-chain descriptions.[8] The PT model illustrates the switch from stick-slip (the typical stop-and-go advance of, say, a chalk stick on a school blackboard) to smooth sliding for decreasing surface corrugation, as well as the passage from large friction at high sliding speeds to vanishingly small friction (thermolubricity) at low speeds.[7,8] In the special case of mismatched lattices, the FK model describes the transition between frictionless (superlubric) motion and pinned frictional sliding (the onset of static friction).

What these models have achieved is to describe how properties such as corrugation, temperature, velocity, and lattice-matching of crystals in contact may influence and determine friction, and thus in principle permit its control, desirable as that is in many practical instances. Despite much theoretical background,[7,8] neither of these models has been really tested experimentally. That is what emulators can do.

The emulators of Ref. 2 and 3 are based on short chains of trapped ionized atoms that, under the influence of an electric field, are forced to slide across a laser-generated optical lattice. These techniques may seem arcane, yet they are accurate and powerful, because parameters such as the temperature, atom velocity and spacing, chain length and the amplitude of the lattices potential, can be flexibly adjusted across a vast range of values.

In this vein, Gangloff and colleagues present an experiment involving mostly a single ion that slides across an optical lattice, a set-up that emulates the PT model to near perfection. Through it they demonstrate that, even at their microKelvin temperature, there is a speed below which the friction between the sliding ion and the lattice vanishes (thermolubricity), as demanded by thermodynamics. Above that speed, the friction reverts to stick-slip and rises above two orders of magnitude with increasing speed. No experiment involving real crystals can span this kind of range. Although much of the authors findings had been known from numerical simulations,[7–9] their emulator is superior – for example it reproduces much better the theoretically expected dependence of friction on velocity.[10]

Emulating the FK model after the PT model would require an infinitely long ion chain in place of a single ion. Bylinskii et al. used instead a short chain of two to six ions sliding across an optical lattice – so is this just a baby step towards that goal? Not quite. By adjusting the distances between the ions, the authors tune the amount of mismatch between the chain and the corrugation of the lattices periodic potential, and produce a dramatic effect on the friction that they measure.

Starting from large friction for perfect chain-lattice matching, Bylinskii and colleagues observe a rapid decrease of the friction as they increase a parameter that controls the amount of chain-lattice mismatch. This trend reflects the evolution from strong pinning friction, a regime where it takes a large force to dislodge and nudge the chain forward, towards better lubricity, the sliding behaviour of two mismatched lattices, whose relative position is energetically more indifferent. Thus for increasing mismatch the system tends towards lubricity; even if not necessarily superlubricity, a regime only reached



FIG. 1. **Ion chains trapped in a lattice.** Bylinskii et al.[2] and Gangloff et al.[3] investigate the frictional forces arising when one or a few ions are forced to slide across optical lattices, where the corrugated intensity potentials is created by counter-propagating lasers. These simple experiments have provided benchmark tests of long-standing frictional models for sliding crystal surfaces.

when the lattice potential intensity falls below the value that demarcates the Aubry transition between frictionless and pinned frictional sliding.[11] It will be interesting in the future to pursue the vestiges of this important transition, known to survive even for very short chains,[4,12] as is the case in this kind of emulators. It is amazing to remark how much already at this stage the study of just a few ions can teach us about the physics of infinite systems.

As always, experiments teach us more than we anticipate. Clear-cut techniques, like the cold-ion emulators reported by these two research papers, provide useful insights into the complexity underlying even the simplest act of friction involving a handful of ions. In this sense, paraphrasing the "more is different" dictum by physicist Philip Warren Anderson,[13] one could here say that sometimes "less in not that different" after all.

This work was mainly supported under the ERC Advanced Grant No. 320796-MODPHYSFRICT.

[1] A. Vanossi, N. Manini, M. Urbakh, S. Zapperi, and E. Tosatti, Rev. Mod. Phys. **85**, 529 (2013).
[2] A. Bylinskii, D. Gangloff, V. Vuletic, Science **348**, 1115-1118 (2015).
[3] D. Gangloff, A. Bylinskii, I. Counts, W. Jhe, V. Vuletic, Nat. Phys. **11**, 915-919 (2015).
[4] A. Benassi, A. Vanossi, E. Tosatti, Nat. Comm. **2**, 236 (2011).
[5] I. Garcia-Mata, O. V. Zhirov, D. L. Shepelyansky, Eur. Phys. J. D **41**, 325 (2007).
[6] T. Pruttivarasin, M. Ramm, I. Talukdar, A. Kreuter, H. Häffner, New J. Phys. **13**, 075012 (2011) .
[7] M. H. Müser, Phys. Rev. B **84**, 125419 (2011).
[8] S. Y. Krylov, J. W. M. Frenken, Phys. Status Solidi B **251**, 711 (2014).
[9] O. M. Braun, Y. Kivshar, "The Frenkel-Kontorova Model: Concepts, Methods and applications" (Springer, Berlin, 1998).
[10] Y. Sang, M. Dubé, M. Grant, Phys. Rev. Lett. **87**, 174301 (2001).
[11] S. Aubry, P. Y. Le Daeron, Physica D **8**, 381 (1983).
[12] S. R. Sharma, B. Bergersen, B. Joos, Phys. Rev. B **29**, 6335 (1984).
[13] P. W. Anderson, Science **177**, 393 (1972).